

Citation for published version:

Hathout, RM & Woodman, TJ 2014, 'NMR diffusion-ordered spectroscopy can explain differences in skin penetration enhancement between microemulsion formulations', *Nanomedicine: Nanotechnology, Biology, and Medicine*, vol. 10, no. 7, pp. 1389-1390. <https://doi.org/10.1016/j.nano.2014.06.006>

DOI:

[10.1016/j.nano.2014.06.006](https://doi.org/10.1016/j.nano.2014.06.006)

Publication date:

2014

Document Version

Peer reviewed version

[Link to publication](#)

University of Bath

Alternative formats

If you require this document in an alternative format, please contact:
openaccess@bath.ac.uk

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.



NMR diffusion-ordered spectroscopy can explain differences in skin penetration enhancement between microemulsion formulations

Rania M. Hathout, PhD^{a,b,*}, Timothy J. Woodman, PhD^b

^aDepartment of Pharmaceutics and Industrial Pharmacy, Faculty of Pharmacy, Ain Shams University, Cairo, Egypt

^bDepartment of Pharmacy and Pharmacology, University of Bath, Bath, UK

Received 26 March 2014; accepted 12 June 2014

Abstract

Changing the formulation variables of microemulsion systems has a significant influence on the resulting transdermal enhancement effect. NMR diffusion-ordered spectroscopy (DOSY) can offer an extremely valuable tool to interpret the differences in the obtained fluxes based on variations in self-diffusions between the drug and its locus domain.

© 2014 Published by Elsevier Inc. All rights reserved.

Key words: Microemulsions; NMR; Diffusion-ordered; Transdermal; Diffusion; Penetration enhancement

Microemulsion systems are usually considered as successful vehicles for transdermal drug delivery.¹ In part owing to the varied component composition, they usually offer excellent skin penetration enhancement. This is usually attributed to three key mechanisms; notably the hydration they impose to the stratum corneum due to their aqueous phase; the solubilizing power they impart to oil soluble drugs and finally the effect of the different surfactants and co-surfactants that are used in their production.^{2,3} In this note, we introduce a new physical insight to interpret differences in penetration enhancement between microemulsion formulations based on studying the self-diffusion of a lipophilic drug *viz* testosterone relative to the diffusion of its residing oil phase. Such information can be readily obtained using diffusion ordered ¹H NMR spectroscopy (DOSY) and subsequently correlation of the values obtained with the testosterone fluxes through porcine skin.

Diffusion-ordered nuclear magnetic resonance (NMR) spectroscopy (DOSY) is a powerful technique that utilizes apparent translational diffusion coefficients in characterizing molecular species.⁴ Several methods exist to extract diffusion coefficients from NMR data, the most visually appealing of which is a *pseudo* two dimensional spectrum where the conventional NMR is displayed in the direct dimension and the diffusion coefficient is depicted in the indirect one. Characteristic peaks for each

component in the mixture may be traced. Recently, many applications have emerged utilizing this valuable technique in drug delivery using microemulsion systems such as: determination of the microemulsion type, detection of aggregates and determination of the drug locus inside the microemulsion.⁵ Also, an attempt has been reported to correlate the self-diffusion coefficient of a certain drug; vinpocetine with its transdermal fluxes, although in this study the diffusion of the residing domain, though important, was neglected, leading to limited results interpretation.⁶

Transdermal fluxes of testosterone incorporated in microemulsion formulations were chosen along the dilution line L20 which starts at (% wt/wt) 20 oleic acid/40 Tween 20®/40 Transcutol® (having different water compositions from 5% to 23%) in the constructed phase diagram of the system oleic acid/Tween 20®/Transcutol®/water, were obtained from Hathout et al., 2010.⁷ Diffusion-ordered spectroscopy (¹H) was performed in order to determine the self-diffusion coefficients of the drug and the oleic acid (the oil phase of the microemulsion and the drug locus).^{7,8} The results were reported by Hathout and Woodman.⁵

Accordingly, testosterone transdermal steady state fluxes (J_{ss}) from the differently prepared microemulsion formulations were correlated with the self-diffusion of testosterone ($D_{\text{testosterone}}$) in these microemulsions normalized with the diffusion of oleic acid ($D_{\text{oleic acid}}$); its locus phase in the microemulsion (Figure 1). A very good correlation ($r = 0.90$) was obtained. It was observed that increasing the water content of the microemulsions had a lower effect on testosterone diffusion compared to oleic acid; the

*Corresponding author.

E-mail address: r_hathout@yahoo.com (R.M. Hathout).




Figure 1. Correlation between testosterone transdermal fluxes across porcine skin and its normalized self-diffusion in the microemulsion system oleic acid/Tween 20®/Transcutol®/water.

ratio $D_{\text{testosterone}}/D_{\text{oleic acid}}$ increased by increasing the microemulsion water percentage. In other words, the difference in mobility between the drug and its domain was increased. Interestingly, this was also accompanied by an increase in testosterone fluxes through the porcine skin. The flux through the skin (drug penetration) usually follows the diffusion (or partitioning) of testosterone to the outer surface of the skin (i.e. $A > B > C$). The slower of these consecutive steps determines the final rate. In concentrated microemulsions, the diffusion step ($A > B$) is slower than the penetration step therefore it becomes the rate-limiting one. It may be expected that, after a certain dilution with water the diffusion becomes faster and the second step becomes rate-limiting. However, in the current study increasing the water content further caused the disappearance of the microemulsion area⁷ so the former case prevailed. We assume that increasing the water content relatively constrains the diffusion of the oily phase (where lipophilic drugs such as testosterone reside) and generates relative diffusion differences between the drug and its domain. This enhances the potential for

the drug to leave its locus and partition to the aqueous phase and the skin surface thereafter or to directly diffuse to the skin surface if the oily domain comes into direct contact with it. Consequently, we encourage potential researchers working on transdermal delivery using microemulsion systems to consider similar correlations in interpreting the microemulsion's composition effects on skin penetration or permeation. Using NMR diffusimetry; a simple but a very effective method for quality control (using low field NMR) and for research (using high field NMR) is a new idea in the field of transdermal delivery research.

References

1. Heuschkel S, Goebel A, Neubert RHH. Microemulsions—modern colloidal carrier for dermal and transdermal drug delivery. *J Pharm Sci* 2008;**97**:603–31.
2. Rhee YS, Huh JY, Park CW, Nam TY, Yoon KR, Chi SC, et al. Effects of vehicles and enhancers on transdermal delivery of clobopride. *Arch Pharm Res* 2007;**30**:1155–61.
3. Trommer H, Neubert RHH. Overcoming the stratum corneum: the modulation of skin penetration—a review. *Skin Pharmacol Physiol* 2006;**19**:106–21.
4. Liu HZ, Wang YJ, Lang YY, Yao HM, Dong Y, Li SM. Bicontinuous cyclosporin A loaded water-AOT/Tween 85-isopropylmyristate microemulsion: structural characterization and dermal pharmacokinetics in vivo. *J Pharm Sci* 2009;**98**:1167–76.
5. Hathout RM, Woodman TJ. Applications of NMR in the characterization of pharmaceutical microemulsions. *J Control Release* 2012;**161**:62–72.
6. Hua L, Weisan P, Jiayu L, Ying Z. Preparation, evaluation, and NMR characterization of vinpocetine microemulsion for transdermal delivery. *Drug Dev Ind Pharm* 2004;**30**:657–66.
7. Hathout RM, Woodman TJ, Mansour S, Mortada ND, Geneidi AS, Guy RH. Microemulsion formulations for the transdermal delivery of testosterone. *Eur J Pharm Sci* 2010;**40**:188–96.
8. Hathout RM, Elshafeey AH. Development and characterization of colloidal soft nano-carriers for transdermal delivery and bioavailability enhancement of an angiotensin II receptor blocker. *Eur J Pharm Biopharm* 2012;**82**:230–40.



Graphical Abstract

NMR diffusion-ordered spectroscopy can explain differences in skin penetration enhancement between microemulsion formulations

Rania M. Hathout^{a,b}, Timothy J. Woodman^b

^aDepartment of Pharmaceutics and Industrial Pharmacy,
Faculty of Pharmacy, Ain Shams University, Cairo, Egypt

^bDepartment of Pharmacy and Pharmacology, University of Bath, Bath, UK

Transdermal flux of testosterone is correlated with the normalized self-diffusion of the drug relative to its locus self-diffusion.

Nanomedicine: Nanotechnology, Biology, and Medicine xxx (2014) xxx

